

VITAMIN D

The past year has produced stunning research findings concerning vitamin D's potential role in preventing and perhaps even treating cancer.

Scientists are examining the use of vitamin D to reduce the risk of no fewer than 17 different types of cancer, ranging from colon, breast, and prostate cancers to ovarian, esophageal, renal, and bladder cancers. Moreover, researchers believe vitamin D may even improve treatment outcomes in people already diagnosed with cancer. *A recent review article estimated that 50,000-70,000 Americans die prematurely from cancer each year due to insufficient intake of vitamin D.*[1]

Emerging research suggests that vitamin D also has applications in promoting bone strength, as well as in mitigating autoimmune conditions such as multiple sclerosis, type I diabetes, and rheumatoid arthritis. Other potential benefits include promoting dental and skin health, relieving depression and helping to prevent stroke, metabolic syndrome, and musculoskeletal pain.

Combating Colorectal Cancer

Vitamin D's effects in reducing cancer risk have been studied most extensively in colorectal cancer, the second leading cause of cancer death in the US.[3] A study in 2005 investigated the relationship between vitamin D intake, serum vitamin D levels, and colorectal cancer risk. Individuals with vitamin D intake of 1000 IU or more daily or with serum vitamin D (25-hydroxyvitamin D) levels of 33 nanograms per milliliter (ng/mL) experienced a 50% lower risk of colorectal cancer. A daily dose of 1000 IU of vitamin D is half the safe upper limit established by the National Academy of Sciences. According to the study authors, prompt public health action is needed to increase daily intake of vitamin D to 1000 IU and to raise serum levels of 25-hydroxyvitamin D. For some individuals, modest sunlight exposure may help achieve these optimal levels.[4]

An epidemiological review conducted in 2005 at Harvard Medical School corroborated vitamin D's protective effects against colorectal cancer and noted that typical dietary intake of 200-400 IU per day is probably too low to confer appreciable benefits. The Harvard study noted that a person's vitamin D status at the time of cancer diagnosis and treatment may influence survival.[5]

In the recent Polyp Prevention Trial, investigators analyzed several dietary factors in relation to the recurrence of adenomatous polyps in the colon.[6] Adenomatous polyps are considered pre-malignant and may therefore be a harbinger of colon cancer. Low intake of calcium and vitamin D was associated with increased risk of recurrence of the pre-malignant polyps. Optimal vitamin D and calcium status may thus be an important preventive strategy against colon cancer.

Protecting the Prostate

Recent clinical trials suggest that vitamin D and its analogs promise to be important therapies for prostate cancer.[7] Experimental evidence indicates that the active form of vitamin D promotes differentiation and inhibits proliferation, invasiveness, and metastasis of human prostate cancer cells.[8,9]

In a study conducted last year, investigators examined the relationship between sun exposure and prostate cancer. Comparing 450 men with advanced prostate cancer with 450 unaffected men, they found that those with a high level of sun exposure had a 50% lower prostate cancer risk than

men with low sun exposure. The researchers believe sunlight helped protect the men against prostate cancer by promoting vitamin D synthesis. Because of the association between sun exposure and certain skin cancers, however, the scientists noted, "increasing vitamin D intake from diet and supplements may be the safest solution to achieve adequate levels of vitamin D." [9]

Another study in 2005 demonstrated a therapeutic role for vitamin D in prostate cancer. Sixteen men who had previously been treated for prostate cancer supplemented with 2000 IU daily of vitamin D. The investigators then monitored prostate-specific antigen (PSA) levels for over two years. PSA is a marker of prostate cancer recurrence or progression. In nine patients, PSA levels decreased or remained unchanged after vitamin D supplementation began. In patients with rising PSA levels, supplementation with vitamin D3 significantly lengthened the PSA doubling time, by an average of 75%. (The rate at which PSA increases or doubles is correlated with disease prognosis, with longer PSA doubling times associated with better outcomes.) These findings indicate that vitamin D may help to slow or prevent disease recurrence or progression in patients who have been treated for prostate cancer. [10]

Vitamin D has also been reported to benefit patients whose prostate cancer has metastasized to the bones. This patient population commonly develops vitamin D deficiency. Supplementing these individuals with vitamin D was found to reduce pain, boost muscle strength, and improve overall quality of life. [11]

Benefits for Breast Health

Several lines of evidence suggest vitamin D may help reduce the incidence of breast cancer. A prospective study published in 2005 examined the relationship of plasma vitamin D metabolites to breast cancer risk in a cohort of women enrolled in the Nurses' Health Study. Blood samples were collected from study participants from 1989 to 1990 and analyzed for the vitamin D metabolites. The study participants were followed until 1996. The researchers then compared blood samples from women who developed breast cancer with samples from cancer-free control subjects. High levels of vitamin D metabolites were associated with a lower risk of breast cancer. For both metabolites, the association was stronger in women aged 60 and older. [12]

In another line of study in 2005, investigators looked at the relationship between dietary vitamin D and calcium intake and breast density as measured by mammography. [13]

Mammographic density is considered a strong risk factor for breast cancer. [14] Dietary intake of vitamin D and calcium was assessed in a group of women, aged 40-60, who had screening mammograms. Women who had a combined daily intake of 100 IU or more of vitamin D combined with 750 mg or more of calcium demonstrated decreased breast density compared to women with lower intakes of the two nutrients. This suggests that adequate consumption of vitamin D and calcium may reduce breast cancer risk by influencing breast tissue architecture. [13]

Improving Lung Cancer Survival

Vitamin D may help people with lung cancer to live longer, according to a study released in 2005. The most common cause of cancer death in American men and women, lung cancer can be challenging to treat effectively. The report found that men with early-stage non-small cell lung cancer who had higher vitamin D indices (based on dietary intake and exposure to sunlight) had improved recurrence-free survival rates compared to men with lower vitamin D intake and sun exposure. [15] Another study conducted in 2005 demonstrated that the active form of vitamin D

inhibited lung cancer metastasis in an animal model of the disease.[16] These findings suggest that implementing vitamin D therapy may be critical to improving survival rates for lung cancer patients.

Additional scientific evidence suggests that optimal vitamin D status may be associated with reduced risk of many other forms of cancer. These include cancers of the bladder, esophagus, stomach, ovary, uterus, cervix, pancreas, larynx, oral cavity, and gall bladder, as well as Hodgkin's and non-Hodgkin's lymphomas.[1]

The Depression Connection

Vitamin D helps the brain produce serotonin, a neurotransmitter critical to emotional health. Vitamin D deficiency can contribute to negative emotions such as depression. Likewise, increased vitamin D consumption elevates mood and promotes a positive outlook.

In particular, supplementation with vitamin D may prevent seasonal depression. The nutrient is produced in the skin during exposure to sunlight. As the days grow shorter and colder in winter, the body may be unable to produce adequate amounts of vitamin D. Individuals living in colder climates where the daylight hours shorten significantly - including those in the Northern United States, Canada, and Northern Europe - are most at risk for vitamin D deficiency and the resulting "winter depression". But clinical research shows that taking extra vitamin D during the winter can improve mood and ward off the wintertime blues.

In one double-blind trial, people received 400-800 IU of vitamin D or placebo for five days during late winter. Those taking vitamin D experienced a significant enhancement in positive mood compared to those taking placebo.[54] Particularly notable is the unusually rapid response produced by vitamin D supplementation. Individuals felt better after taking vitamin D for only five days. [54]

Preventing Multiple Sclerosis

Multiple sclerosis is an immune-mediated inflammatory and neurodegenerative condition of the central nervous system. Its symptoms include weakness, visual problems, and impaired coordination. Although its causes remain unknown, scientists suspect that multiple sclerosis may represent an autoimmune condition.

Investigators have established a strong link between multiple sclerosis incidence and geographic location, noting that areas with abundant sun exposure or plentiful dietary fish intake experience reduced risk. Multiple sclerosis occurs more often in people who lived in northern areas of Europe and North America during childhood, and less often in people who live closer to the equator.[23] Individuals appear to retain the level of risk associated with the area in which they lived until age 15, even if they moved to a different area later in life.[23] In Switzerland, multiple sclerosis rates are higher at low altitudes and lower at high altitudes, where UV light is more intense. In Norway, multiple sclerosis rates are higher inland, but much lower near the coast, where vitamin D₃-rich fish is consumed regularly.[24] Vitamin D, obtained through both sun exposure and diet, may be the factor responsible for the link between geography and multiple sclerosis risk.

Evidence suggests that vitamin D supplementation may decrease the lifetime risk of multiple sclerosis in women. Experimental data suggest the white matter of the brain that multiple sclerosis affects contains vitamin D receptors, and inadequate vitamin D during early development may predispose these cells to an early demise.[25]

Effects on Type I Diabetes

Some scientists believe that type I diabetes may be an autoimmune condition in which insulin-producing pancreatic beta cells are destroyed. Evidence from animal experiments and human observational studies suggests that vitamin D may help prevent type I diabetes, perhaps by acting as an immune system modulator.[27]

Human studies likewise suggest that vitamin D may have a protective effect against type I diabetes. In a large-scale investigation, more than 12,000 pregnant women in Finland enrolled in a trial studying the relationship between vitamin D intake and type I diabetes in infants. After one year, children who supplemented with the suggested study dose of vitamin D (2000 IU per day) had a much lower risk of type I diabetes than children who did not supplement.[29]

Lowering Gingivitis Risk

High blood levels of a vitamin D metabolite are associated with a decreased risk of the gum disease gingivitis, according to a recent report from the American Journal of Clinical Nutrition. Researchers at Boston University analyzed data from 6,700 nonsmokers, aged 13-90+, from the National Health and Nutrition Examination Survey.[30]

Participants with the highest blood levels of Vitamin D were the least likely to display signs of gingivitis. In fact, the association between vitamin D levels and gingivitis incidence appeared to be linear over the entire range of blood levels. This association was similar even in relation to other factors such as gender, ethnic groups, and age.[30]

The scientists noted that vitamin D may reduce susceptibility to gingivitis by exerting anti-inflammatory effects, and postulated that gingivitis may provide a useful clinical model for further investigation into the anti-inflammatory effects of vitamin D.[30]

Promoting Bone Health

One of vitamin D's greatest contributions to health is promoting strong, healthy bones. Vitamin D deficiency is associated with skeletal diseases characterized by weak bones, such as rickets in children and osteomalacia and osteoporosis in adults.[22]

Vitamin D combined with calcium supplementation is widely known to help decrease postmenopausal bone loss and prevent osteoporosis.[31,32] Furthermore, vitamin D combined with calcium can help decrease the risk of hip and non-vertebral fractures.[32]

Osteoporosis prevention may optimally begin early in life. In a retrospective study, investigators compared prepubescent females who received oral vitamin D in infancy to those who did not. Girls who received vitamin D had significantly increased bone mineral density compared to those who did not receive the vitamin.[33]

While commonly thought of as a female disease, osteoporosis affects men as well. Osteoporosis and associated fractures are increasingly prevalent in men, and mortality rates following major fractures are higher in men than in women.[34] As with women, osteoporosis prevention in men should begin in youth and continue in adulthood, using vitamin D, calcium supplementation, and physical activity.

Alleviating Musculoskeletal Pain

Low levels of vitamin D are associated with persistent, non-specific musculoskeletal pain, according to investigators at the University of Minnesota Medical School. Researchers conducted a cross-sectional study of 150 patients, aged 10-65, who presented to a primary care

clinic over the course of two years with the complaint of persistent, non-specific musculoskeletal pain. Serum vitamin D levels were analyzed to assess vitamin D status.[36]

Ninety-three percent of the patients demonstrated deficient levels of vitamin D, and 28% were considered severely deficient. Five patients had vitamin D levels that were too low to detect. *The research team concluded that all patients-regardless of gender or age-with chronic, non-specific musculoskeletal pain are at high risk of suffering from unrecognized vitamin D deficiency.*

Preventing Falls in the Elderly

In elderly adults, falls occur frequently and are associated with significant morbidity and mortality.[38] Research suggests that vitamin D may help prevent these dangerous falls.

Cross-sectional studies have shown that elderly adults with higher serum levels of vitamin D demonstrate a lower number of falls, as well as increased muscle strength.[39] One meta-analysis found that vitamin D supplementation helped reduce the risk of falling by more than 20%.[38] A randomized, controlled trial found that three months of supplementing with 1200 mg of calcium and 800 IU of vitamin D3 daily reduced the risk of falling by 49% in adults in a long-term geriatric care facility.[39]

NEW APPLICATIONS FOR VITAMIN D

In recent months, scientists have discovered that vitamin D may have applications in preventing and managing numerous health conditions, including:

Rheumatoid arthritis. This inflammatory condition causes swelling, immobility, and joint pain. Recent reports indicate that vitamin D deficiency in adults is associated with an elevated risk of developing rheumatoid arthritis.[40] Vitamin D may also be a crucial therapeutic tool for helping individuals receiving corticosteroid treatment for rheumatoid arthritis to maintain optimal bone mass.[41]

Hypertension. Investigators note that mounting evidence suggests that vitamin D deficiency may increase the risk of hypertension.[42] A recent study identified a possible mechanism by which vitamin D may support healthy blood pressure. Researchers at the University of Chicago found that analogs of vitamin D inhibit the expression of renin, a kidney-produced hormone involved in the pathogenesis of hypertension, in both laboratory and animal studies. This suggests that vitamin D could affect blood pressure by inhibiting renin, which helps regulate blood pressure.[43]

Skin disorders. Some investigators believe that based on the distribution of vitamin D receptors throughout the body, vitamin D may have potential therapeutic applications in managing numerous skin disorders, including actinic keratosis, seborrheic dermatitis, and photoaging.[44] One of the most troublesome and difficult-to-treat skin conditions is psoriasis, which is marked by patches of itchy, flaky skin. Both ultraviolet B radiation and topical application of the active form of vitamin D have been found to benefit patients who suffer from psoriasis. Scientists believe that vitamin D's antiproliferative and immune-regulating activities may be responsible for these benefits.[45,46]

Metabolic Syndrome. A precursor to type II diabetes, metabolic syndrome is associated with an increased risk of heart disease. A recent study analyzing data from more than 10,000 participants found that increased dietary intake of vitamin D was associated with a decreased risk of developing metabolic syndrome. Calcium intake also exerted a protective effect against metabolic syndrome in this study of middle-aged and older women.[47]

Averting stroke. A recent study examined how dietary intake and serum levels of vitamins and minerals influenced stroke risk in elderly subjects. More than 700 men and women were followed for up to 10 years. Low intake of vitamin D and low serum levels of its active form significantly predicted an increased risk for stroke, even after adjusting for factors such as age, gender, and smoking.[48]

Preventing tooth loss. Abundant intake of vitamin D is associated with a decreased risk of tooth loss, according to a recent study conducted in Japan. Researchers found that individuals in their seventies with relatively higher intakes of certain nutrients-including protein, vitamin D, and B vitamins-demonstrated a reduced incidence of missing teeth compared to people who consumed less of these nutrients.[49]

Supporting lung health. People with higher levels of vitamin D in the bloodstream demonstrate better lung function than their counterparts with lower vitamin D levels, according to a recent report. Researchers examined 14,000 participants and found that people with higher vitamin D levels could inhale and exhale substantially more air, as measured by forced vital capacity and forced expiratory volume. Though further studies are needed, these findings suggest that vitamin D might benefit smokers, asthmatics, and others with compromised respiratory function.[50]

THE SUNSHINE DEBATE

Because vitamin D is fat soluble, it can be stored in the adipose tissue of the body, presumably for long-term access. Much debate on vitamin D has focused on the need for dietary supplementation versus the body's endogenous (internal) manufacture of the vitamin from sunlight exposure.

For those living in climates with greatly reduced angles of sunlight (the northern and southern parts of the globe) or where sunlight itself is rare, the need for supplementation is unchallenged. The human body is designed to obtain vitamin D from exposure to sunlight, with only brief exposure providing roughly 80-90% of the body's vitamin D stores.[17] Exposure of the entire body to sunlight may produce approximately 10,000 IU of vitamin D a day.[18] To prevent the accumulation of toxic levels of vitamin D, the body naturally limits the amount of vitamin D it synthesizes from sunlight.[19]

For years, health advocates have suggested that sun exposure may contribute to cancer risk and sunscreen should be used for all sun exposure greater than 15 minutes. New evidence suggests, however, that vitamin D can in fact protect against several forms of cancer. While sunscreen may help protect against the deadliest skin cancer-melanoma-its effect of limiting vitamin D production could lead to a greater incidence of other cancers.[20] That does not mean that sunscreens should not be used, as they can significantly protect against premature skin aging (and skin cancers). What this does tell us, however, is that in one way or another, it is critical that we obtain optimal levels of vitamin D.

Cancer occurs more frequently in dark-skinned people, the obese, and regions with limited exposure to ultraviolet B radiation from sunlight. Each of these factors is associated with low blood levels of vitamin D. Furthermore, cancer survival rates are lower when the diagnosis occurs in months of lower sunlight levels, suggesting a protective role of vitamin D. Studies suggest that vitamin D protects against numerous forms of cancer, including widely prevalent cancers such as those affecting the colon, prostate, breast, and lung.[1,20]

Safety

Vitamin D is typically well tolerated in adults at doses up to 2000 IU daily, with some research indicating that even higher levels up to 10,000 IU daily may be used safely without adverse effects.[22,53] Excess vitamin D can lead to symptoms such as nausea, vomiting, poor appetite, constipation, and weakness.[22] Vitamin D is contraindicated in individuals with elevated blood calcium levels or hypercalcemia.[21] Individuals with kidney disease and people who use digoxin or other cardiac glycoside drugs should consult a physician before using supplemental vitamin D.[21]

Conclusion

Once considered little more than a compound that promotes healthy bones, vitamin D is now recognized as an important weapon in the fight against cancer. Its many other health-promoting effects include protecting muscle strength and modulating autoimmune disease. Optimizing vitamin D status through supplementation and prudent sun exposure should be a cornerstone of every health maintenance program.

Recommended Vitamin D supplements:

<u>Rating</u>	<u>Product</u>	<u>Website</u>
★★★★★	Metagenics- ISO D3 2000	www.drpez.com
★★★★	Vitamin D3 1000	www.lef.org
★★★	Vitamin D 700	www.gnc.com

To order from drpez.com:

Go to www.drpez.com and click on **Patient Gateway** in the upper right-hand corner. Then click on **Register Now** in the upper left-hand corner and follow the prompts. This will allow you access to the Metagenics product gateway.

Metagenics provides the highest quality supplements available today. Their Vitamin D is certified to assure purity and predicted safety. Metagenics sells their products only through health care professionals familiar with their appropriate use.

References

1. Grant WB, Holick MF. Benefits and requirements of vitamin D for optimal health: a review. *Altern Med Rev.* 2005 Jun; 10(2):94-111.
2. van den Bemd GJ, Chang GT. Vitamin D and vitamin D analogs in cancer treatment. *Curr Drug Targets.* 2002 Feb;3(1):85-94.
3. Available at: <http://www.cdc.gov/cancer/colorectl/colorect.htm>. Accessed November 17, 2005.
4. Gorham ED, Garland CF, Garland FC, et al. Vitamin D and prevention of colorectal cancer. *J Steroid Biochem Mol Biol.* 2005 Oct;97(1-2):179-94.
5. Giovannucci E. The epidemiology of vitamin D and colorectal cancer: recent findings. *Curr Opin Gastroenterol.* 2006 Jan;22(1):24-9.
6. Hartman TJ, Albert PS, Snyder K, et al. The association of calcium and vitamin D with risk of colorectal adenomas. *J Nutr.* 2005 Feb;135(2):252-9.
7. Vijayakumar S, Mehta RR, Boerner PS, Packianathan S, Mehta RG. Clinical trials involving vitamin D analogs in prostate cancer. *Cancer J.* 2005 Sep-Oct;11(5):362-73.
8. Lou YR, Qiao S, Talonpoika R, Syvala H, Tuohimaa P. The role of Vitamin D3 metabolism in prostate cancer. *J Steroid Biochem Mol Biol.* 2004 Nov;92(4):317-25.
9. John EM, Schwartz GG, Koo J, Van Den BD, Ingles SA. Sun exposure, vitamin D receptor gene polymorphisms, and risk of advanced prostate cancer. *Cancer Res.* 2005 Jun 15;65(12):5470-9.
10. Woo TC, Choo R, Jamieson M, Chander S, Vieth R. Pilot study: potential role of vitamin D (Cholecalciferol) in patients with PSA relapse after definitive therapy. *Nutr Cancer.* 2005;51(1):32-6.
11. van Veldhuizen PJ, Taylor SA, Williamson S, Drees BM. Treatment of vitamin D deficiency in patients with metastatic prostate cancer may improve bone pain and muscle strength. *J Urol.* 2000 Jan;163(1):187-90.
12. Bertone-Johnson ER, Chen WY, Holick MF, et al. Plasma 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev.* 2005 Aug;14(8):1991-7.

AGE-MANAGEMENT MEDICINE ♦ FOR A LIFETIME OF HEALTH

13. Berube S, Diorio C, Verhoek-Ofstedahl W, Brisson J. Vitamin D, calcium, and mammographic breast densities. *Cancer Epidemiol Biomarkers Prev.* 2004 Sep;13(9):1466-72.
14. Cerhan JR, Sellers TA, Janney CA, et al. Prenatal and perinatal correlates of adult mammographic breast density. *Cancer Epidemiol Biomarkers Prev.* 2005 Jun;14(6):1502-8.
15. Zhou W, Suk R, Liu G, et al. Vitamin D is associated with improved survival in early-stage non-small cell lung cancer patients. *Cancer Epidemiol Biomarkers Prev.* 2005 Oct;14(10):2303-9.
16. Nakagawa K, Kawaura A, Kato S, Takeda E, Okano T. 1 alpha,25-Dihydroxyvitamin D(3) is a preventive factor in the metastasis of lung cancer. *Carcinogenesis.* 2005 Feb;26(2):429-40.
17. Shearer MJ. The roles of vitamins D and K in bone health and osteoporosis prevention. *Proc Nutr Soc.* 1997 Nov;56(3):915-37.
18. Gesensway D. Vitamin D. *Ann Intern Med.* 2000 Aug 15;133(4):319-20.
19. Holick MF. Sunlight and vitamin D: both good for cardiovascular health. *J Gen Intern Med.* 2002 Sep;17(9):733-5.
20. Giovannucci E. The epidemiology of vitamin D and cancer incidence and mortality: a review (United States). *Cancer Causes Control.* 2005 Mar;16(2):83-95.
21. Available at: http://www.pdrhealth.com/drug_info/nmdrugprofiles/nutsupdrugs/vit_0265.shtml. Accessed November 17, 2005.
22. Available at: <http://ods.od.nih.gov/factsheets/vitaminD.asp>. Accessed November 17, 2005.
23. Franklin GM, Nelson L. Environmental risk factors in multiple sclerosis: causes, triggers, and patient autonomy. *Neurology.* 2003 Oct 28;61(8):1032-4.
24. Hayes CE, Cantorna MT, DeLuca HF. Vitamin D and multiple sclerosis. *Proc Soc Exp Biol Med.* 1997 Oct;216(1):21-7.
25. Chaudhuri A. Why we should offer routine vitamin D supplementation in pregnancy and childhood to prevent multiple sclerosis. *Med Hypotheses.* 2005;64(3):608-18.
26. VanAmerongen BM, Dijkstra CD, Lips P, Polman CH. Multiple sclerosis and vitamin D: an update. *Eur J Clin Nutr.* 2004 Aug;58(8):1095-109.
27. Hyponen E. Micronutrients and the risk of type 1 diabetes: vitamin D, vitamin E, and nicotinamide. *Nutr Rev.* 2004 Sep;62(9):340-7.
28. Gysemans CA, Cardozo AK, Callewaert H, et al. 1,25-Dihydroxyvitamin D3 modulates expression of chemokines and cytokines in pancreatic islets: implications for prevention of diabetes in nonobese diabetic mice. *Endocrinology.* 2005 Apr;146(4):1956-64.
29. Hyponen E, Laara E, Reunanen A, Jarvelin MR, Virtanen SM. Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet.* 2001 Nov 3;358(9292):1500-3.
30. Dietrich T, Nunn M, Wason-Hughes B, Bischoff-Ferrari HA. Association between serum concentrations of 25-hydroxyvitamin D and gingival inflammation. *Am J Clin Nutr.* 2005 Sep;82(3):575-80.
31. Anon. ACOG practice bulletin. Clinical management guidelines for obstetrician-gynecologists. Number 50, January 2003. *Obstet Gynecol.* 2004 Jan;103(1):203-16.
32. Bischoff-Ferrari HA, Willett WC, Wong JB, et al. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA.* 2005 May 11;293(18):2257-64.
33. Zamora SA, Rizzoli R, Belli DC, Slosman DO, Bonjour JP. Vitamin D supplementation during infancy is associated with higher bone mineral mass in prepubertal girls. *J Clin Endocrinol Metab.* 1999 Dec;84(12):4541-4.
34. Faraawi R. Osteoporosis in men. *Saudi Med J.* 2005 Apr;26(4):519-23.
35. Heaney RP. Advances in therapy for osteoporosis. *Clin Med Res.* 2003 Apr;1(2):93-9.
36. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc.* 2003 Dec;78(12):1463-70.
37. Holick MF. Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart disease, and osteoporosis. *Am J Clin Nutr.* 2004 Mar;79(3):362-71.
38. Bischoff-Ferrari HA, Wason-Hughes B, Willett WC, et al. Effect of Vitamin D on falls: a meta-analysis. *JAMA.* 2004 Apr 28;291(16):1999-2006.
39. Bischoff HA, Stahelin HB, Dick W, et al. Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. *J Bone Miner Res.* 2003 Feb;18(2):343-51.
40. Holick MF. Vitamin D: important for prevention of osteoporosis, cardiovascular heart disease, type 1 diabetes, autoimmune diseases, and some cancers. *South Med J.* 2005 Oct;98(10):1024-7.
41. Miggiano GA, Gagliardi L. Diet, nutrition, and rheumatoid arthritis. *Clin Ter.* 2005 May-Jun;156(3):115-23.
42. Holick MF. The vitamin D epidemic and its health consequences. *J Nutr.* 2005 Nov;135(11):2739S-48S.
43. Qiao G, Kong J, Uskokovic M, Li YC. Analogs of 1alpha,25-dihydroxyvitamin D(3) as novel inhibitors of renin biosynthesis. *J Steroid Biochem Mol Biol.* 2005 Jun;96(1):59-66.
44. Nagpal S, Na S, Rathnachalam R. Noncalcemic actions of vitamin D receptor ligands. *Endocr Rev.* 2005 Aug;26(5):662-87.
45. Lehmann B. The vitamin D3 pathway in human skin and its role for regulation of biological processes. *Photochem Photobiol.* 2005 Feb 1.
46. Wolters M. Diet and psoriasis: experimental data and clinical evidence. *Br J Dermatol.* 2005 Oct;153(4):706-14.
47. Liu S, Sonh Y, Ford ES, Manson JE, Buring JE, Ridker PM. Dietary calcium, vitamin D, and the prevalence of metabolic syndrome in middle-aged and older US women. *Diabetes Care.* 2005 Dec;28(12):2926-32.
48. Marniemi J, Alanen E, Impivaara O, et al. Dietary and serum vitamins and minerals as predictors of myocardial infarction and stroke in elderly subjects. *Nutr Metab Cardiovasc Dis.* 2005 Jun;15(3):188-97.
49. Yoshihara A, Watanabe R, Nishimuta M, Hanada N, Miyazaki H. The relationship between dietary intake and the number of teeth in elderly Japanese subjects. *Gerodontology.* 2005 Dec;22(4):211-8.
50. Available at: <http://today.reuters.co.uk/news/newsArticleSearch.aspx?storyID=259205+12-Dec+2005+RTRS&srch=vitamin+d>. Accessed December 13, 2005.
51. Available at: http://today.reuters.co.uk/news/newsArticle.aspx?type=healthNews&storyID=2005-11-17T050400Z_01_CC718202_RTRIDST_0_HEALTH-VITAMIN-DC.XML. Accessed November 17, 2005.
52. Steingrimsdottir L, Gunnarsson O, Indridason OS, Franzson L, Sigurdsson G. Relationship between serum parathyroid hormone levels, vitamin D sufficiency, and calcium intake. *JAMA.* 2005 Nov 9;294(18):2336-41.
53. Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr.* 1999 May;69(5):842-56.
54. Lansdowne ATG, Provost SC. Vitamin D3 enhances mood in healthy subjects during winter. *Psychopharmacology* 1998 135:319-23.